

ANOPHELES DIRUS SPECIES E: CHROMOSOMAL AND CROSSING EVIDENCE FOR ANOTHER MEMBER OF THE DIRUS COMPLEX¹

Y. SAWADIPANICH,² V. BAIMAI² AND B. A. HARRISON³

ABSTRACT. Cytogenetic and crossing data provide strong evidence for the existence of another species, *dirus* E in southwestern India, within the Dirus Complex of *Anopheles*. These findings are in accord with unpublished morphological observations. Our data suggest a significant genetic divergence between species E and its close relatives, *An. dirus* A, B and C in Thailand. These data also suggest that *dirus* E is an incipient sibling species of its geographically nearest relative, *dirus* D, and that it seemingly co-evolved through the process of allopatric speciation.

INTRODUCTION

The Dirus Complex of *Anopheles* subgenus *Cellia* consists of at least 7 morphologically similar species which occur from southwestern India across the Southeast Asian mainland to Con Son Island, Vietnam, Hainan Island, China and Taiwan (Peyton and Ramalingam 1988, Baimai 1988, Peyton 1990). These include 3 described species: *An. dirus* Peyton and Harrison, 1979 (species A), *An. nemophilous* Peyton and Ramalingam, 1988 (previously species F of Baimai et al. 1988a) and *An. takasagoensis* Morishita, 1946 (Peyton and Harrison 1980). Currently undescribed members of the complex are provisionally designated as species B, C, D (Baimai et al. 1987) and species E (Tewari et al. 1987). On August 24, 1981, a laboratory colony of the little-known *dirus* member from southwestern India was established by one of us (BAH) at the Department of Medical Entomology, Armed Forces Research Institute of Medical Sciences (AFRIMS), Bangkok, from eggs provided upon request by Hari R. Bhat, National Institute of Virology, Pune, India. Preliminary cross-mating studies indicated that the southwest Indian *dirus* member was different from *An. balabacensis* Baisas, from Sabah, Malaysia (V. Baimai, unpublished data). For this reason, cytogenetic and crossing studies were performed to determine whether species E was really a member of the Dirus Complex. This paper presents the results of these investigations.

MATERIALS AND METHODS

Use of the designation "Dirus Complex" is taken from Peyton and Ramalingam (1988) and

Peyton (1990). Although *An. dirus* is the correct name for what previously was designated species A, we have chosen to continue the use of "species A" for convenience in our crossing tables.

The original colony of *dirus* E was established from pooled egg-batches of 8 females collected in Shimoga District, Karnataka, southwestern India, during August 1981. This colony was maintained in the laboratory at 26°C by the artificial mating method (Ow Yang et al. 1963). All combinations of crosses (at least 10 pairs for each cross) between this colony and the laboratory stocks of 4 species of the Dirus Complex were performed by forced matings. The species strains used in this study were species A from Phet Buri (TL 33, 1983), species B from Phatthalung (PT 59, 1985), species C from Nakhon Si Thammarat (SC 28, 1984) and species D from Ranong (CP 25, 1983). These colonies were derived from isofemale lines and maintained in the insectary at Mahidol University.

Mitotic and salivary gland polytene chromosomes were prepared from fourth-instar larvae using the modified methods of Baimai et al. (1981). The male larva (XY), has a relatively thin polytene X chromosome, as compared with the normal thickness of a X chromosome of a female larva (XX). The fertility of F₁ hybrid progeny was determined by egg hatch success from the back- and self-crossing experiments, and from examination of the testes of F₁ hybrid males. Also, the degree of asynapsis in salivary gland polytene chromosomes of F₁ hybrids was taken to reflect the degree of genetic incompatibility.

RESULTS AND DISCUSSION

The mitotic karyotype of species E is very similar to that of *dirus* D (Baimai et al. 1987), particularly the sex chromosomes (cf. species A and B, Baimai et al. 1981). Giemsa staining revealed that the X chromosome of species E has a short telocentric shape with 2 separate dark bands of intercalary heterochromatin in the euchromatic section (Figs. 1-3). Only a small block of heterochromatin was observed at the centromeric region of the X chromosome of

¹ The views of the authors do not purport to reflect the position of the Department of the Army or the Department of Defense.

² Department of Biology, Faculty of Science, Mahidol University, Rama VI Road, Bangkok 10400, Thailand.

³ Walter Reed Biosystematics Unit, Department of Entomology, Walter Reed Army Institute of Research, Washington, DC 20307-5100.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE SEP 1990		2. REPORT TYPE		3. DATES COVERED 00-00-1990 to 00-00-1990	
4. TITLE AND SUBTITLE Anopheles Dirus Species E: Chromosomal and Crossing Evidence for Another Member of the Dirus Complex				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Department of Entomology, Washington, DC, 20307				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			




Fig. 1-4. Mitotic karyotype from larval neuroblast cells: 1, 2, female and male, respectively, of *Anopheles dirus* species E; 3, 4, F₁ hybrid female and male, respectively, from cross between *dirus* A female and *dirus* E male ($\times 670$) (an arrow indicates a major block of heterochromatin).

species E, compared with that of species A (Fig. 3). The Y chromosome is small and rod-shaped and is mostly heterochromatin (Figs. 2, 4). This chromosome exhibits a large dotlike appearance in some preparations. Each of the autosomes have small blocks of pericentromeric heterochromatin similar to those of species A.

In the crossing studies, extensive asynapsis (over 90%) was observed in salivary gland polytene chromosomes of F₁ hybrid larvae derived from the crosses between species E and species A, B and C (Fig. 5) and heterozygous inversions were observed in the X chromosome and chromosome arms 2L and 2R in all cases (see Baimai et al. 1988b). However, F₁ hybrid larvae from the crosses between species E and D exhibited only small sections of asynapsis in the salivary gland polytene chromosomes, particularly at the very tips of chromosome arms 2L, 2R and 3R (Fig. 6), and nearly complete synapsis along the X chromosome except at the very tip. These observations suggest that a fixed simple inversion is likely to exist in the X chromosome of

species E similar to that of species D. These data also suggest that species E possesses floating inversions in at least 2 autosome arms.

The fertility results of cross-mating experiments between species E and the 4 species of the complex from Thailand are summarized in Table 1. All combinations of crosses produced viable F₁ hybrids of both sexes. However, when F₁ hybrid progeny were backcrossed to the parental species only a very few progeny (larvae which subsequently died) were obtained in 3 crosses, and F₁ hybrids self-crossed failed to produce viable F₂ eggs (Table 2). The microscopic examination of testes of F₁ males from all of the crosses revealed that they were abnormal in size and shape (Figs. 7, 8) and contained either non-motile sperm or no sperm.

Crosses between species A and E produced a large number of eggs in both directions. The cross between females of species A and males of species E yielded relatively low percentages of hatched eggs (37.2%), but larval survival was comparable with that of the reciprocal cross




Fig. 5, 6. Larval polytene chromosomes from F₁ hybrid female larvae resulting from crosses between *Anopheles dirus* A female and *dirus* E male ($\times 260$) and *dirus* D female and *dirus* E male ($\times 130$), respectively (arrows indicate paracentric inversions). Figures 7, 8. Abnormal testes and accessory glands of F₁ hybrid males from the respective cross-matings in Figs. 5 and 6.

Table 1. Results of cross-mating of *Anopheles dirus* A, B, C and D from Thailand and species E from southwestern India.

Crosses		No. of ovipositions hatched (total)	Mean no. of eggs per oviposition	% eggs hatched	% larval survival	% F ₁ adults emerged	% F ₁ males*
female × male							
E	A	12 (15)	84.7	75.9 (772/1016)	73.8 (570)	89.1 (508)	48.6
A	E	10 (11)	91.2	37.2 (339/912)	65.8 (223)	53.3 (119)	58.8
E	B	4 (14)	71.3	2.5 (7/285)	71.4 (5)	60.0 (3)	33.3
B	E	10 (11)	81.9	71.3 (584/819)	22.1 (129)	70.1 (93)	52.7
E	C	13 (15)	96.6	56.8 (713/1256)	68.7 (490)	98.8 (484)	46.5
C	E	10 (11)	101.7	84.8 (862/1017)	83.8 (722)	93.9 (678)	53.7
E	D	4 (10)	81.0	51.5 (167/324)	49.1 (82)	81.7 (67)	56.7
D	E	5 (7)	103.6	76.1 (394/518)	43.2 (170)	94.7 (161)	57.8

* All F₁ male hybrids were sterile.

Table 2. Back- and self-crossing experiments with the F₁ hybrids from the crosses listed in Table 1.

Crosses		No. of ovipositions	% eggs hatched* (no.)
female × male		(total pair matings)	
F ₁ (EA)	E	0 (5)	—
F ₁ (EA)	A	1 (5)	16.7 (10/60)
E	F ₁ (EA)	0 (5)	—
A	F ₁ (EA)	0 (5)	—
F ₁ (EA)	F ₁ (EA)	3 (5)	0 (0/228)
F ₁ (AE)	E	0 (5)	—
F ₁ (AE)	A	9 (11)	36.0 (180/500)
E	F ₁ (AE)	0 (5)	—
A	F ₁ (AE)	0 (5)	—
F ₁ (AE)	F ₁ (AE)	0 (10)	—
F ₁ (BE)	E	0 (5)	—
F ₁ (BE)	B	0 (5)	—
E	F ₁ (BE)	0 (5)	—
B	F ₁ (BE)	0 (5)	—
F ₁ (BE)	F ₁ (BE)	0 (10)	—
F ₁ (CE)	E	0 (5)	—
F ₁ (CE)	C	0 (5)	—
E	F ₁ (CE)	0 (5)	—
C	F ₁ (CE)	0 (5)	—
F ₁ (CE)	F ₁ (CE)	0 (5)	—
F ₁ (EC)	E	0 (5)	—
F ₁ (EC)	C	3 (5)	7.5 (15/200)
E	F ₁ (EC)	0 (5)	—
C	F ₁ (EC)	0 (5)	—
F ₁ (EC)	F ₁ (EC)	3 (5)	0 (0/80)
F ₁ (DE)	E	0 (5)	—
F ₁ (DE)	D	0 (5)	—
E	F ₁ (DE)	0 (5)	—
D	F ₁ (DE)	0 (5)	—
F ₁ (ED)	F ₁ (DE)	0 (5)	—

* All larvae died before pupation.

(65.8 vs. 73.8%). The opposite result was obtained for the crosses between species C and E. Crosses between females of species E and males of species B were the least successful; only 2.5% of the eggs hatched and few larvae survived to become adults. The reciprocal cross gave a high

percentage of hatched eggs (71.3%) but a low percentage of surviving larvae (22.1%). A greater degree of genetic compatibility was observed in the crosses between species D and E. Crosses in both directions produced fairly high percentages of hatched eggs (51.5 and 76.1%) and moderate survival (49.1 and 43.2%). These data suggest that there may be different degrees of genetic incompatibility at different stages of development of the F₁ hybrids derived from the crosses between species E and *dirus* A, B, C and D.

The above cytogenetic and crossing evidence clearly indicates that species E from southwestern India represents a distinct species, and is the seventh member recognized so far within the *Dirus* Complex. These findings are in complete accord with ongoing morphological studies (E. L. Peyton, unpublished data). Species E has been designated *dirus* E by Tewari et al. (1987), Peyton and Ramalingam (1988), Bhat (1988) and Peyton (1990). Our cytogenetic data suggest that *dirus* E is more closely related to *dirus* D than to the other members of the complex studied here. This is supported by data from recent studies of the geographic distribution of *dirus* D (Baimai et al. 1988c), apparently the predominant species throughout The Union of Myanmar [Burma] and Bangladesh. Populations of *dirus* E in southwestern India are the most western members of the *Dirus* Complex (see the comments of Bhat 1988, regarding a record of *bala-bacensis* from Kasauli in the western Himalayas), and are currently isolated from populations of *dirus* D. Thus, it seems probable that *dirus* E could have arisen from an ancestral stock of *dirus* D through the process of allopatric speciation. If this is the case, *dirus* E is probably the only representative of the *Dirus* Complex in southwestern India. The recognition of *dirus* E has proven to be of considerable importance in understanding the evolution of this medically important complex of Oriental *Anopheles*.

ACKNOWLEDGMENTS

We are very grateful to R. E. Harbach and E. L. Peyton for commenting on this manuscript, K. Vejsanit for assistance in preparing the illustrations and L. Panthusiri for maintaining the AFRIMS colony of *dirus* E. This work was supported by the UNDP/World Bank/WHO Special Program for Research and Training in Tropical Diseases.

REFERENCES CITED

- Baimai, V. 1988. Population cytogenetics of the malaria vector *Anopheles leucosphyrus* group. Southeast Asian J. Trop. Med. Public Health 19:667-680.
- Baimai, V., B. A. Harrison and L. Somchit. 1981. Karyotype differentiation of 3 anopheline taxa in the Balabacensis Complex of Southeast Asia (Diptera: Culicidae). Genetica 57:81-86.
- Baimai, V., R. G. Andre, B. A. Harrison, U. Kijchalao and L. Panthusiri. 1987. Crossing and chromosomal evidence for two additional sibling species within the taxon *Anopheles dirus* Peyton and Harrison (Diptera: Culicidae) in Thailand. Proc. Entomol. Soc. Wash. 89:157-166.
- Baimai, V., R. E. Harbach and U. Kijchalao. 1988a. Cytogenetic evidence for a fifth species within the taxon *Anopheles dirus* in Thailand. J. Am. Mosq. Control Assoc. 4:333-338.
- Baimai, V., A. Poopittayasataporn and U. Kijchalao. 1988b. Cytological differences and chromosomal rearrangements in four members of the *Anopheles dirus* complex (Diptera: Culicidae). Genome 30:372-379.
- Baimai, V., Myat Myat Thu, Myo Paing and N. P. Maheswary. 1988c. Distribution and chromosomal polymorphism of the malaria vector *Anopheles dirus* species D. Southeast Asian J. Trop. Med. Public Health 19:661-665.
- Bhat, H. R. 1988. A note on *Anopheles dirus* Peyton and Harrison, 1979 [*An. balabacensis* (sensu lato) Baisas, 1936] in India. Indian J. Malariol. 25:103-105.
- Morishita, K. 1946. A new variety of Formosan *Anopheles*—*Anopheles leucosphyrus* var. *takasagoensis* var. nov. J. Formosan Med. Assoc. 45:21, 65-70.
- Ow Yang, C. K., F. L. Sta Maria and R. H. Wharton. 1963. Maintenance of laboratory colony of *Anopheles maculatus* Theobald by artificial mating. Mosq. News 23:34-35.
- Peyton, E. L. 1990. A new classification for the Leucosphyrus Group of *Anopheles* (*Cellia*). Mosq. Syst. (1989) 21:197-205.
- Peyton, E. L. and B. A. Harrison. 1979. *Anopheles* (*Cellia*) *dirus*, a new species of the Leucosphyrus Group from Thailand (Diptera: Culicidae). Mosq. Syst. 11:40-52.
- Peyton, E. L. and B. A. Harrison. 1980. *Anopheles* (*Cellia*) *takasagoensis* Morishita 1946, an additional species in the Balabacensis Complex of Southeast Asia (Diptera: Culicidae). Mosq. Syst. 12:335-347.
- Peyton, E. L. and S. Ramalingam. 1988. *Anopheles* (*Cellia*) *nemophilous*, a new species of the Leucosphyrus Group from Peninsular Malaysia and Thailand (Diptera: Culicidae). Mosq. Syst. 20:272-299.
- Tewari, S. C., J. Hiriyan and R. Reuben. 1987. Survey of the anopheline fauna of the Western Ghats in Tamil Nadu, India. Indian J. Malariol. 24:21-28.